

Cholesterol medication list

Lipid-lowering drug classes, statin intensity and safety, and a patient therapy record

Page 1 is the class reference. Page 2 sets statin intensity by dose and the interactions and monitoring that go with it. Page 3 is the patient record, including the trial history payers ask for at prior authorization.

1. LIPID-LOWERING DRUG CLASSES

CLASS AND MECHANISM	GENERIC (BRAND)	DOSE RANGE	LDL-C REDUCTION	INDICATION
Statins Block HMG-CoA reductase, so the liver makes less cholesterol and pulls more LDL from the blood	Atorvastatin (Lipitor), rosuvastatin (Crestor), simvastatin (Zocor), pravastatin (Pravachol), lovastatin (Altoprev), fluvastatin (Lescol XL), pitavastatin (Livalo)	See page 2. Doses run from simvastatin 10 mg to atorvastatin 80 mg and rosuvastatin 40 mg	18–63% by drug and dose	First line. Clinical ASCVD, LDL-C 190 mg/dL or above, diabetes at ages 40–75, and primary prevention by calculated risk
Cholesterol absorption inhibitor Blocks NPC1L1 uptake in the small intestine	Ezetimibe (Zetia)	10 mg once daily	13–20% alone; about 20–25% added to a statin	The usual first add-on when a statin alone misses the LDL-C target, and an option in statin intolerance
PCSK9 monoclonal antibodies Bind PCSK9 so LDL receptors are recycled rather than degraded	Alirocumab (Praluent), evolocumab (Repatha)	Alirocumab 75–150 mg SC every 2 weeks, or 300 mg every 4 weeks. Evolocumab 140 mg SC every 2 weeks, or 420 mg monthly	50–60%	Clinical ASCVD or familial hypercholesterolemia when maximally tolerated statin and ezetimibe fall short
siRNA against PCSK9 Silences hepatic PCSK9 mRNA before the protein is made	Inclisiran (Leqvio)	284 mg SC at baseline, again at 3 months, then every 6 months	About 50%	Adjunct in ASCVD or familial hypercholesterolemia. Twice-yearly dosing suits a patient who will not self-inject
ATP citrate lyase inhibitor Blocks cholesterol synthesis upstream of HMG-CoA reductase; activated only in the liver, not muscle	Bempedoic acid (Nexletol); with ezetimibe (Nexlizet)	180 mg once daily; 180/10 mg as the combination	17–18% alone; about 38% with ezetimibe	Statin intolerance, and as an adjunct. Carries a cardiovascular risk-reduction indication. Lowers uric acid in some patients but raises it in others
Bile acid sequestrants Bind bile acids in the gut, forcing the liver to use cholesterol to replace them	Cholestyramine (Prevalite), colestevlam (Welchol), colestipol (Colestid)	Cholestyramine 4–16 g/day; colestevlam 3.75 g/day; colestipol 2–16 g/day	15–30%	Adjunct where a statin is not tolerated. Raises triglycerides, so avoid when they are already high
Fibrates PPAR-alpha agonists that clear triglyceride-rich particles	Fenofibrate (Tricor, Trilipix), gemfibrozil (Lopid)	Fenofibrate 48–145 mg daily; gemfibrozil 600 mg twice daily	Variable, and LDL-C can rise	Severe hypertriglyceridemia, mainly to prevent pancreatitis. Lowers triglycerides 30–50%
Omega-3 fatty acids Reduce hepatic VLDL assembly and secretion	Icosapent ethyl (Vascepa), omega-3-acid ethyl esters (Lovaza)	Icosapent ethyl 2 g twice daily; omega-3-acid ethyl esters 4 g daily	Minimal	Triglycerides of 500 mg/dL or above. Icosapent ethyl also reduces cardiovascular events on top of a statin
Niacin Reduces hepatic triglyceride synthesis and VLDL release	Extended-release niacin (Niaspan), immediate-release (Niacor)	500–2,000 mg at bedtime, titrated slowly	5–20%	Not recommended for routine cardiovascular risk reduction. Two large trials found no added benefit on top of a statin, with more harm
Combination products Two mechanisms in one tablet	Ezetimibe / simvastatin (Vytorin), bempedoic acid / ezetimibe (Nexlizet), amlodipine / atorvastatin (Caduet)	Vytorin 10/10 to 10/40 mg; Nexlizet 180/10 mg	As for the components combined	Fewer tablets where a patient already needs both agents. Check the statin component against the interactions on page 2

Homozygous familial hypercholesterolemia has its own agents, which sit outside the ten classes above: lomitapide (Juxtapid), an MTP inhibitor, and evinacumab (Evkeeza), an ANGPTL3 antibody given by infusion. Both are restricted to that diagnosis.

2. STATIN INTENSITY BY DOSE

INTENSITY	DRUG AND DAILY DOSE
High LDL-C down 50% or more	Atorvastatin 40–80 mg • Rosuvastatin 20–40 mg
Moderate LDL-C down 30–49%	Atorvastatin 10–20 mg • Rosuvastatin 5–10 mg • Simvastatin 20–40 mg • Pravastatin 40–80 mg • Lovastatin 40–80 mg • Fluvastatin XL 80 mg • Fluvastatin 40 mg twice daily • Pitavastatin 1–4 mg
Low LDL-C down less than 30%	Simvastatin 10 mg • Pravastatin 10–20 mg • Lovastatin 20 mg • Fluvastatin 20–40 mg

Three traps worth checking before you write the dose. **Rosuvastatin 5 mg and atorvastatin 10 mg are moderate intensity, not high** — the drug name does not set the band, the dose does. **Pitavastatin never reaches high intensity** at any approved dose. And **simvastatin 80 mg** is restricted to patients who have already taken it for 12 months without muscle injury. Bands as defined by the ACC/AHA dyslipidemia guideline; the 2026 revision keeps the 2018 thresholds.

KEY INTERACTIONS TO SCREEN BEFORE PRESCRIBING

COMBINATION	WHAT HAPPENS	WHAT TO DO
Simvastatin or lovastatin + strong CYP3A4 inhibitors (clarithromycin, erythromycin, itraconazole, ketoconazole, posaconazole, voriconazole, HIV protease inhibitors, cobicistat, nefazodone)	Statin levels rise sharply, and with them the risk of rhabdomyolysis	Contraindicated. Switch the antibiotic, or hold the statin for the course
Simvastatin + verapamil, diltiazem or dronedarone	Moderate rise in simvastatin exposure	Cap simvastatin at 10 mg daily. With amiodarone, amlodipine or ranolazine, cap it at 20 mg
Any statin + gemfibrozil	Myopathy risk rises well above that of either drug alone	Avoid the pairing. Use fenofibrate when a fibrate is needed alongside a statin
Rosuvastatin + cyclosporine or gemfibrozil	Transporter inhibition raises rosuvastatin levels	Cap rosuvastatin at 5 mg with cyclosporine, and at 10 mg with gemfibrozil
Bempedoic acid + simvastatin or pravastatin	Simvastatin exposure roughly doubles; pravastatin rises about 1.5-fold	Avoid simvastatin above 20 mg and pravastatin above 40 mg
Bile acid sequestrants + levothyroxine, digoxin, warfarin, fat-soluble vitamins	The resin binds the other drug in the gut and absorption drops	Separate doses by at least 4 hours. Colesevelam interacts least
Fibrates + warfarin	INR rises	Check INR within a week of starting or stopping, and adjust the warfarin dose
Ezetimibe + cyclosporine	Both drug levels rise	Monitor cyclosporine levels and liver function
Statins + azithromycin	Negligible CYP3A4 inhibition	No change needed. This is the macrolide to reach for when a statin must continue

MONITORING

Lipid panel — 4 to 12 weeks. Repeat 4 to 12 weeks after starting therapy or changing a dose, to confirm both the response and that the patient is taking it. Then every 3 to 12 months once stable.

ALT — at baseline only. Measure hepatic transaminases before the first statin dose. Routine periodic liver testing is no longer recommended, so repeat only if the patient develops symptoms such as unusual fatigue, abdominal pain, jaundice or dark urine.

Creatine kinase — only if symptoms appear. Do not order it routinely. Measure CK when a patient reports muscle pain, tenderness, weakness or cramping, and take a baseline only in patients already at raised risk of muscle events.

Worth adding by agent. Uric acid on bempedoic acid, which also carries a tendon rupture warning. Triglycerides on a bile acid sequestrant, which pushes them up. HbA1c or fasting glucose on a statin, which modestly raises the chance of new-onset diabetes without outweighing the cardiovascular benefit.

3. PATIENT LIPID THERAPY RECORD

Patient name: _____

Date of birth: _____ mm / dd / yyyy

MRN: _____

Date of review: _____ mm / dd / yyyy

Risk category: _____

LDL-C goal (mg/dL): _____

Lipid results

DATE	TOTAL CHOLESTEROL	LDL-C	HDL-C	TRIGLYCERIDES	NON-HDL-C

All values in mg/dL. Non-HDL-C is total cholesterol minus HDL-C, and needs no fasting sample.

Current agents

DRUG (GENERIC / BRAND)	DOSE AND FREQUENCY	START DATE	INDICATION OR TARGET

Agents tried and stopped

Record every agent the patient has come off, and why. Payers ask for this at prior authorization: approval for a PCSK9 inhibitor or inclisiran usually turns on documented trials of statins at an adequate dose, with the dates and the reason each one ended.

AGENT	DOSE REACHED	DATES (FROM – TO)	REASON STOPPED	DOCUMENTED BY

Reasons to choose from: muscle symptoms with or without raised CK, raised transaminases, inadequate LDL-C response at the maximum tolerated dose, drug interaction, patient preference, cost or coverage.

Next review

Next lab due: _____ mm / dd / yyyy

Tests requested: _____

Clinician signature

Printed name and role

Date

This template is a clinical reference and a documentation aid. It does not replace FDA prescribing information, current guideline recommendations, or your own clinical judgment.