

Pressure Ulcer Nursing Care Plan

Pressure Ulcer Nursing Care Plan

Patient Information

Full Name

Date of Birth

Gender

Patient ID

Contact Number

Email Address

Potential use of medical devices that increase the risk of PI development

- ☐ Respiratory devices
- ☐ Orthopedic devices
- ☐ Urinary or fecal-collecting devices
- ☐ Repositioning devices
- ☐ Drains
- ☐ Tubes
- ☐ Intravenous catheters and lines
- ☐ Restraints
- ☐ Stockings
- ☐ Bandaging

Impaired Skin Integrity - Assessment

Impaired Skin Integrity - Intervention

Impaired Skin Integrity - Notes

Risk for Infection - Assessment

Risk for Infection - Intervention

Risk for Infection - Notes

Risk for Impaired Peripheral Tissue Perfusion - Assessment

Risk for Impaired Peripheral Tissue Perfusion - Intervention

Risk for Impaired Peripheral Tissue Perfusion - Notes

Stage 1 pressure injury - Location

Stage 1 pressure injury - Intervention

Stage 2 pressure injury - Location

Stage 2 pressure injury - Intervention

Stage 3 pressure injury - Location

Stage 3 pressure injury - Intervention

Stage 4 pressure injury - Location

Stage 4 pressure injury - Intervention

Unstageable pressure injury - Location

Unstageable pressure injury - Intervention

Deep pressure injury - Location

Deep pressure injury - Intervention

Physician's Notes and Recommendations

Physician's Signature

Date